

References

1. Richards MJ, Edwards JR, Culver DH, Gaynes RP. Nosocomial infections in combined medical-surgical intensive care units in the United States. *Infect Control Epidemiol* 2000;21:510-5.
2. Centers for Disease Control and Prevention. Guidelines for the prevention of intravascular catheter-related infections. *MMWR* 2002;51(RR-10):1-18.
3. Berenholtz SM, Pronovost PJ, Lipsett PA, et al. Eliminating catheter-related bloodstream infections in the intensive care unit. *Crit Care Med* 2004;32:2014-20.
4. Trautmann M, Zauser B, Wiedeck H, Buttenschön K, Marre R. Bacterial colonization and endotoxin contamination of intravenous infusion fluids. *J Hosp Infect* 1997;37:225-36.
5. Worthington T, Tebbs S, Moss H, Bevan V, Kilburn J, Elliott TS. Are contaminated flush solutions an overlooked source for catheter-related sepsis? *J Hosp Infect* 2001;49:81-3.
6. Calop J, Bosson JL, Croizé J, Laurent PE. Maintenance of peripheral and central intravenous infusion devices by 0.9% sodium chloride with or without heparin as a potential source of catheter microbial contamination. *J Hosp Infect* 2000;46:161-2.
7. Canadian ICU Collaboration. Safer Healthcare Now. Canadian Patient Safety Institute. Available at <http://www.canadianpatientsafetyinstitute.ca>. Accessed July 12, 2007.
8. Central Line Bundle. Available at www.ihl.org. Accessed April 18, 2007.
9. Centers for Disease Control and Prevention. Reduction in central line-associated bloodstream infections among patients in intensive care units—Pennsylvania, April 2001–March 2005. *MMWR Morb Mortal Wkly Rep* 2005;54:1013-16.
10. Field K, McFarlane C, Cheng AC, et al. Incidence of catheter-related bloodstream infection among patients with a needleless mechanical valve-based intravenous connector in an Australian hematology-oncology unit. *Infect Control Hosp Epidemiol* 2007;28:610-3.
11. Rupp ME, Sholtz LA, Jourdan DR, et al. Outbreak of bloodstream infection temporally associated with the use of an intravascular needleless valve. *Clin Infect Dis* 2007;44:1408-14.

12. Maragakis LL, Bradley KL, Song X, et al. Increased catheter-related bloodstream infection rates after the introduction of a new mechanical valve intravenous access port. *Infect Control Hosp Epidemiol* 2006;27:67-70.
13. Hall KK, Geffers C, Gianetta E, Farr BM. Outbreak of bloodstream infections temporally associated with a new needleless IV infusion system. In: Program and abstracts of the 14th Annual Scientific Meeting of the Society for Healthcare Epidemiology of America; April 19, 2004; Philadelphia, PA. Abstract 285.
14. Karchmer TB, Cook EM, Palavecino E, Ohl CA, Sheretz RJ. Needleless valve ports may be associated with a high rate of catheter-related bloodstream infection. 15th Annual Scientific Meeting of the Society for Healthcare Epidemiology of America; April 9–12, 2005; Los Angeles, CA.
15. Salgado CV, Chinnes L, Paczesny TH, Cantey R. Increased rate of catheter-related bloodstream infection associated with use of a needleless mechanical valve device at a long-term acute care hospital. *Infect Control Hosp Epidemiol* 2007;28:684-8.
16. Jarvis W, Sheretz RJ, Perl T, Bradley K, Giannetta E. Increased central venous catheter-associated bloodstream infection rates temporally associated with changing from a split-septum (SS) to a luer-access mechanical valve needleless device: a nationwide outbreak? *Am J Infect Control* 2005;33:E14-15. Abstract 54459.
17. Adams D, Karpanen T, Worthington T, Lambert P, Elliott TSJ. Infection risk associated with a closed luer access device. *J Hosp Infect* 2006;62:353-7.
18. Menyhay SZ, Maki DG. Disinfection of needleless catheter connectors and access ports with alcohol may not prevent microbial entry: The promise of a novel antiseptic-barrier cap. *Infect Control Hosp Epidemiol* 2006; 27:23-27.
19. Hanchett M. Needleless connectors and bacteremia: Is there a relationship? *Infection Control Today* 2005.

20. Goldman DA, Pier GB. Pathogenesis of infections related to intravascular catheterization. Clin Microbiol Rev 1993;6:176-92.